



Fishes of Alto Jacuí sub-basin: a poorly studied sub-basin of northwestern Rio Grande do Sul, Brazil

Karine O. Bonato* and **Clarice B. Fialho**

Universidade Federal do Rio Grande do Sul, Instituto de Biociências, Departamento de Zoologia, Programa de Pós-Graduação em Biologia Animal, CEP 91501-970, Porto Alegre, RS, Brazil

* Corresponding author. E-mail: kakabio2005@yahoo.com.br

Abstract: The streams in the state of Rio Grande do Sul (mainly in the Alto Jacuí sub-basin belonging to the Laguna dos Patos system) have scarce information about their ichthyofauna. Thereby for providing information about stream species, the purpose of the present study was to inventory the ichthyofauna of the streams of the Alto Jacuí sub-basin, located in northwestern state of Rio Grande do Sul. The samples were taken bimonthly from June 2012 to June 2013 using electrofishing technique in 10 streams. A total of 13,247 specimens were collected belonging to 42 species, 10 families and six orders. We report the occurrence of five new species that have not yet been described by researchers.

Key words: stream; Alto Jacuí sub-basin; Laguna dos Patos system; ichthyofauna inventory

INTRODUCTION

Fish are considered the most diverse group of vertebrates (Lowe-MacConnell 1999), with an estimated richness of 32,900 species (Froese and Pauly 2014). By December 2013, Pelayo-Villamil et al. (2014) had found 14,782 described species of fish that occur only in freshwater. Although there is a lack of a complementary information, current estimates of the ichthyofauna in the Neotropical region are that there are about 6,000 to 8,000 species, totaling 13% of the vertebrate biodiversity in aquatic ecosystems worldwide, with Brazilian continental waters showing 21% of global diversity (Reis et al. 2003; Agostinho et al. 2005).

There is still a lack of knowledge of fish richness, mainly in South America, Africa and Asia, which is due to a lack of sampling and databasing (Pelayo-Villamil et al. 2014). Brazil has the largest river networks in the world (Galves et al. 2009); however, many Brazilian basins and sub-basins have not yet been sampled (Agostinho et al. 2005), or there exists little information about

their fish fauna, especially with medium-sized and small water bodies such as streams (Castro 1999). According to Langeani et al. (2007) streams are the environments that have the highest number of new species still to be discovered. But the small size of streams and headwater environments makes these places more susceptible to anthropogenic action and they may experience significant change in their population structure, leading to the disappearance of the most sensitive species (Galves et al. 2009). This situation makes it difficult to understand ecological, biological and biogeographical processes (Barletta et al. 2010).

Although the situation has been changing in the last two decades with an increase of studies (biological, ecological and systematic studies) in streams, there is a lack of knowledge in some regions such as in southern Brazil. Most studies in the state of Rio Grande do Sul refer to large rivers, lagoons or estuarine areas (Fialho et al. 1998; Garcia and Vieira 2001; Bastos 2002; Garcia et al. 2003; Majolo 2005; Maltchik et al. 2005; Dufech and Fialho 2006; Garcia et al. 2006; Petry and Schulz 2006; Ribeiro and Köhler 2007; Dufech and Fialho 2009; Flores-Lopes et al. 2010; Saccol-Pereira and Fialho 2010). The only stream environment studies from Rio Grande do Sul are: Tagliani (1994), Becker (2002), Bozzeti and Schulz (2004), Vilella et al. (2004), Hirschmann (2009), Silva (2009), Winkler-Sosinski et al. (2009), Costa and Schulz (2010) and Volcan et al. (2011).

Little information is available about sampling and studies of the ichthyofauna in streams for the Jacuí river basin and no studies of the upper region of the basin, called the Alto Jacuí sub-basin. Malabarba (1989) showed a list of freshwater fish present in the Laguna dos Patos system and cited species found in Jacuí River and its tributaries. Alves and Fontoura (2009) identified the distributive pattern of migratory fish of the Jacuí River basin, but the data were obtained through interviews, collections, literature and technical studies (EIA-RIMA

Estudo e Relatório de Impacto Ambiental) developed in the study region. Additionally there are some taxonomic reviews and descriptions of new species that are distributed on this drainage (Ottoni and Cheffe 2009; Menezes and Ribeiro 2010; Carvalho and Reis 2011).

We emphasize that to understand the ecological mechanisms in these little-explored environments we must use many tools, including ichthyofaunal studies. Streams are highly heterogeneous environments (Winemiller et al. 2008) and this allows for the establishment of numerous species of fish. Further, more studies of streams in south Brazil are necessary because some basins are not as well explored as the Alto Jacuí sub-basin. Therefore, the aim of this study is to inventory and provide more information about distribution and species richness of ichthyofauna in the Alto Jacuí sub-basin located in northwestern Rio Grande do Sul.

MATERIALS AND METHODS

Study site

The Alto Jacuí sub-basin belongs to the large Laguna dos Patos system and is located in the state of Rio Grande do Sul in the northwestern Middle Plateau and Central Depression region. The Alto Jacuí has its headwaters located in the municipality of Passo Fundo and occupies an area of 16,062 km² with its rivers flowing into the Lago Guaíba (COAJU 2009). The basin's vegetation consists of Seasonal Deciduous Forest and some areas of

Subtropical Ombrophilous Forest. The economy is based on agriculture (soybeans, corn, wheat and rice) and livestock. The basin is drained by the Jacuí, Jacuí-Mirim, Jacuizinho, Caixões, Ivaí and Soturno rivers (SEMA 2010). The Jacuí River is the main tributary of the basin and it is responsible for 85% of the waters forming the Lago Guaíba (FEPAM 2011).

Thus, this study was conducted in 10 streams (Figures 1 and 2–11) in northwestern Rio Grande do Sul, which corresponds to the Alto Jacuí sub-basin (Table 1). All streams flow into the Jacuí River, which is one of the main tributaries to the Laguna dos Patos system.

Data collection

Fish samples were collected with authorization number 34940 from register number 3196382 from Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio). This study was approved by the Ethics Committee on Animal Use of the Universidade Federal do Rio Grande do Sul (permit number 24434) and was conducted in accordance with protocols in their ethical and methodological aspects for the use of fish.

The fish were collected in June, August, October and December 2012; February, April and June 2013. Each sampling event lasted four days. For the sampling, we used electrofishing with three stages of 30 min each, in stretches of 50 m per sampling stream. After sampling, fish were euthanized with 10% eugenol (Vidal et al. 2008; Lucena et al. 2013a), fixed in 10% formalin




Figure 1. Sampling streams in the Alto Jacuí sub-basin. For stream code see Table 1.

and then transferred to 70% alcohol for conservation. The taxonomic identification was carried out in the laboratory using Rodriguez and Reis (2008), Bertaco and

Lucena (2010), Ferrer and Malabarba (2013), Lucena et al. (2013b), Lucena and Soares (2016) and additional literature cited herein. Classification and nomenclature follows Reis et al. (2003), with additional changes made by Thomaz et al. (2015) for Characidae. The nomenclature for Cichlidae followed the new classification of bony fishes proposed by Betancur et al. (2013) that include this family in the order Cichliformes. The voucher specimens were deposited in the fish collection of the Departamento de Zoologia at Universidade Federal do Rio Grande do Sul (UFRGS; Table 2).

RESULTS

A total of 13,247 specimens belonging to 42 species, ten families and six orders (Table 2 and Figures 12–32) were collected. The most significant orders were Characiformes (36%), Cichliformes (24%) and Siluriformes (14%), with 15, 10 and six species, respectively. Cyprinodontiformes was represented by two species, and both Gymnotiformes and Synbranchiformes were represented by only one species. The predominant families were Characidae (12 species, 29%), Loricariidae (nine species, 21%), Cichlidae (seven species, 17%) and Heptapteridae (four species, 10%), followed by Crenuchidae, Poeciliidae and Trychomycteridae with two species (5% each). Erythrinidae, Pimelodidae, Gymnotidae and Synbranchidae showed one species, corresponding to 2% each of the total richness.

Five species are identified only to genus level and correspond to undescribed species: *Australoheros* sp. (Rícan and Kullander 2008), *Bryconamericus* sp. b (Silva 1998), *Heptapterus* sp. (Bockmann 1998), *Ituglanis* sp. (J. Ferrer, personal communication) and *Bryconamericus* sp. a which also seem to be a new species but, could not be described because it may just be a variation (as color and body shape) of *Bryconamericus iheringii*. According to Bonato and Ferrer (2013), the individuals of *Phalloceros spiloura* Lucinda, 2008 collected in the Alto Jacuí sub-basin during the present study represent the first record of this species to the Laguna dos Patos system.

The highest species richness was found in RP, RT, RC and RQ with 28 species, 27 species, 25 species and 21 species, respectively. RM and RSC showed the lowest species richness with only 15 and 14 sample species.




Figures 2–11. General view of the sampled streams in the Alto Jacuí sub-basin, Rio Grande do Sul, Brazil: 2: AA; 3: RP; 4: RD; 5: RC; 6: RJ; 7: RM; 8: RQ; 9: RSC; 10: RT; 11: RV. For stream code see Table 1.

Table 1. Geographic coordinates, elevation and localization of the sampled streams and their respective codes in the Alto Jacuí sub-basin.

Stream	Code	Geographic coordinates	Elevation (m)	Locality
Caixões River	RC	29°01'54.4" S, 052°49'25.1" W	420	Guanxuma, Espumoso
Jacuzinho River	RJ	28°58'02.9" S, 052°47'20.3" W	513	Depósito, Espumoso
Morcego River	RM	28°53'55.0" S, 052°49'05.6" W	461	São Domingos, Espumoso
Turvo River	RT	28°43'47.0" S, 052°47'40.4" W	351	Santo Antônio, Espumoso
Quati River	RQ	28°38'31.8" S, 052°37'07.9" W	439	Mormaço
Santa Clara River	RSC	28°44'30.1" S, 053°13'03.0" W	439	Santa Clara, XV de Novembro
Valoroso Stream	RV	28°41'32.0" S, 052°51'41.5" W	376	Teutônia, Tapera
Divinéia Stream	RD	28°42'16.7" S, 052°52'25.9" W	350	Teutônia, Tapera
Paz Stream	RP	28°42'57.3" S, 052°50'41.7" W	378	Vila Paz, Tapera
Angico Stream	AA	28°39'17.9" S, 052°54'31.1" W	368	São Rafael, Tapera

Table 2. List of fish species collected at each sampled stream in the Alto Jacuí sub-basin. See Table 1 for stream names. Asterisk indicates the endemic species to Laguna dos Patos system.

Taxa	Streams										Voucher (UFRGS)
	RC	RJ	RM	RV	RD	AA	RSC	RQ	RP	RT	
CHARACIFORMES											
Characidae											
<i>Astyanax lacustris</i> (Lütken, 1875)	X			X	X	X		X	X	X	19977
<i>Astyanax laticeps</i> (Cope, 1894)					X		X			X	19327
<i>Astyanax obscurus</i> (Hensel, 1870) *	X						X		X		19329
<i>Astyanax procerus</i> Lucena, Castro & Bertaco, 2013 *	X	X	X	X	X	X	X	X	X	X	19323
<i>Astyanax xiru</i> Lucena, Castro & Bertaco, 2013	X		X		X	X			X	X	19325
<i>Bryconamericus iheringii</i> (Boulenger, 1887)	X	X	X	X	X	X	X	X	X	X	19974
<i>Bryconamericus</i> sp. a	X	X									19975
<i>Bryconamericus</i> sp. b *	X										19980
<i>Diapoma alburnus</i> (Hensel, 1870)	X	X	X					X	X	X	19976
<i>Diapoma dicropotamicus</i> (Malabarba & Weitzman, 2003) *			X					X			19952
<i>Oligosarcus jacuiensis</i> Menezes & Ribeiro, 2010	X	X			X	X	X	X	X	X	19978
<i>Oligosarcus jenynsii</i> (Günther, 1864)				X		X		X	X	X	19979
Crenuchidae											
<i>Characidium orientale</i> Buckup & Reis, 1997	X										19962
<i>Characidium pterostictum</i> Gomes, 1947	X	X	X	X	X	X	X	X	X	X	19973
Erythrinidae											
<i>Hoplias malabaricus</i> (Bloch, 1794)		X	X			X			X	X	19961
SILURIFORMES											
Heptapteridae											
<i>Rhamdia quelen</i> (Quoy & Gaimard, 1824)	X	X	X	X		X	X	X	X	X	19265
<i>Rhamdella eriarcha</i> (Eigenmann & Eigenmann, 1888)	X										19951
<i>Heptapterus mustelinus</i> (Valenciennes, 1835)	X	X					X				19967
<i>Heptapterus</i> sp.	X	X	X	X	X	X	X	X	X	X	19266
Trichomycteridae											
<i>Ituglanis</i> sp.									X		19949
<i>Trichomycterus poikilos</i> Ferrer & Malabarba, 2013 *	X	X	X	X	X	X	X	X	X	X	19267
Loricariidae											
<i>Ancistrus brevipinnis</i> (Regan, 1904)	X	X	X	X	X	X	X	X	X	X	19984
<i>Eurycheilichthys limulus</i> Reis & Schaefer, 1998	X	X	X	X	X	X	X	X	X	X	19985
<i>Hemiancistrus punctulatus</i> Cardoso & Malabarba, 1999 *	X	X		X	X	X		X	X	X	19986
<i>Hisonotus armatus</i> Carvalho, Lehmann, Pereira & Reis, 2008 *	X										19957
<i>Hisonotus brunneus</i> Carvalho & Reis, 2011 *		X						X			19959
<i>Hypostomus commersoni</i> (Valenciennes, 1836)					X				X		19958
<i>Rineloricaria baliola</i> Rodriguez & Reis, 2008		X	X	X	X	X	X	X	X	X	19982
<i>Rineloricaria cadeae</i> (Hensel, 1868) *				X		X		X	X	X	19983
<i>Rineloricaria microlepidogaster</i> (Regan, 1904)	X	X	X								19981
Pimelodidae											
<i>Pimelodus pintado</i> Azpelicueta, Lundberg & Loureiro, 2008									X	X	19960
GYMNOTIFORMES											
Gymnotidae											
<i>Gymnotus</i> aff. <i>carapo</i> Linnaeus, 1758	X							X	X	X	19966
CICHLIFORMES											
Cichlidae											
<i>Australoheros</i> sp.	X	X	X					X		X	19968
<i>Crenicichla lepidota</i> Heckel, 1840				X	X				X	X	19969
<i>Crenicichla punctata</i> Hensel, 1870	X			X	X			X	X	X	19972
<i>Geophagus brasiliensis</i> (Quoy & Gaimard, 1824)				X			X		X	X	19965
<i>Gymnogeophagus gymnogenys</i> (Hensel, 1870)									X	X	19964
<i>Gymnogeophagus labiatus</i> (Hensel, 1870)	X										19954
<i>Gymnogeophagus rhabdotus</i> (Hensel, 1870)								X		X	19955
CYPRINODONTIFORMES											
Poeciliidae											
<i>Phalloceros caudimaculatus</i> (Hensel, 1868)				X		X			X	X	19963
<i>Phalloceros spiloura</i> Lucinda, 2008				X	X	X					17827
SYNBRANCHIFORMES											
Synbranchidae											
<i>Synbranchus marmoratus</i> Bloch, 1795									X		19953
Total of Species (n= 42)	25	18	15	18	17	18	14	21	28	27	



Figures 12–32. Some species of fishes found in Alto Jacuí sub-basin, Rio Grande do Sul. Photos by L.R. Malabarba. **12:** *Astyanax xiru*, **13:** *Astyanax procerus*, **14:** *Bryconamericus* sp. b (UFRGS 17931), **15:** *Bryconamericus iheringii* (UFRGS 17992), **16:** *Australoheros* sp.(UFRGS 17989), **17:** *Trichomycterus poikilos*, **18:** *Heptapterus* sp. (UFRGS 17933), **19:** *Heptapterus mustelinus*, **20:** *Rhamdella eriarcha* (UFRGS 17984), **21:** *Characidium pterostictum* (UFRGS 17826), **22:** *Characidium orientale*, **23:** *Eurycheilichthys limulus* (UFRGS 17986), **24:** *Rineloricaria baliola* (UFRGS 17991), **25:** *Rineloricaria cadeae*, **26:** *Hyposotomus commersoni*, **27:** *Crenicichla punctata* (UFRGS 17990), **28:** *Gymnogeophagus gymnogenys*, **29:** *Rhamdia quelen*, **30:** *Gymnotus* aff. *carapo* (UFRGS 17989), **31:** *Astyanax laticeps*, **32:** *Astyanax obscurus*.

DISCUSSION

According to Pelayo-Villamil et al. (2014) an average of 240.2 species of fishes were described per year in the last ten years worldwide. The five new species uncovered by this inventory (*Australoheros* sp., *Heptapterus* sp., *Bryconamericus* sp. a and b, and *Ituglanis* sp.) support the importance of this type of study. In addition, inventories are important in extending the distributional range of some species, such as *Phalloceros spiloura* that was previously only known from the coastal drainages of states of Rio Grande do Sul and Santa Catarina, Iguaçu and Uruguay river basins, and as part of this study, was found in the Alto Jacuí sub-basin representing a new record for the Laguna dos Patos system (Bonato and Ferrer 2013).

Malabarba (1989) registered 25 species of the 42 species sampled in this study to Laguna dos Patos system. The most recent literature indicates a total of 160 species to the Laguna dos Patos system (Malabarba et al. 2009) including 35 species that were new species and yet not described in 2009. Of these 35 species listed by Malabarba et al. (2009), we have sampled five species that were described in recent years (*Oligosarcus jacuiensis* Menezes & Ribeiro, 2010; *Hisonotus brunneus* Carvalho & Reis, 2011; *Astyanax procerus* Lucena, Castro & Bertaco, 2013; *Astyanax xiru* Lucena, Castro & Bertaco, 2013; *Trychomycterus poikilos* Ferrer & Malabarba, 2013), indicating that a representative amount of the ichthyofauna of the upper Jacuí River was described in recent years. There are no comparable studies for the Alto Jacuí sub-basin. We can only make comparisons with other basins belonging to the Laguna dos Patos system. For stream environments Bozzeti and Schulz (2004) found 57 species in the Gravataí and Sinos sub-basins, Hirschmann (2009) found 55 species in the Forqueta sub-basin (Taquari-Antas basin), and Becker et al. (2013) found 119 species for the Taquari-Antas basin but in respect to the last study, the high number of captured species is likely due to their larger sampling of 519 sites.

The number of species found in this study is lower compared to those cited by other studies mainly because it was conducted in streams from a headwater region and many of the streams (of lower species richness) are first-order. The highest occurrence of the orders Characiformes, Cichliformes and Siluriformes in studies is also well documented for the Laguna dos Patos system and for the Neotropical region (Castro 1999; Garcia et al. 2003; Buckup et al. 2007; Lévêque et al. 2008; Costa and Schulz 2010). Headwater streams do not have an exclusive fish fauna but, species that form populations residing in streams and also that occur in larger bodies of water with different characteristics (Castro 1999). The fish fauna of streams is based on small species and according to Castro (1999) it seems to be the only

general pattern with real diagnostic value for stream environments. In this study, the streams with lower species richness are the first-order streams, which have a habitat of lower complexity as RD, RV and AA streams (see Table 1 for stream codes). This situation is expected in accordance with the River Continuum Theory (Vannote et al. 1980). Thus, the larger streams, with greater width between banks, areas with and without shading, and more heterogeneous environment showed the highest species richness (Ferreira and Casatti 2006; Suárez and Petrere-Junior 2005) as occurred in the RT, RP and RC streams.

Despite the fact that we did not evaluate the degree of anthropic influences in the sampled streams, all streams sampled here showed some kind of human interference. Most streams are very close to agricultural areas with the presence of dairy cattle or pig livestock. In stream RP there was a considerable amount of waste coming from homes and sometimes we found dead animals within the stream. Probably the residents of the region slaughter animals for their own consumption and discard the remains of the animal in the river. However, this stream had considerable marginal vegetation and a heterogeneous environment with changing pools and areas of rapids, which led to the high amount of richness observed.

Due to a lack of data for streams of the studied sub-basin it is difficult to say that the number of species found is representative of the streams belonging to Laguna dos Patos system. The checklist showed 42 species representing 26% of the species mentioned for the Laguna dos Patos system. This study is an important record for the region of the Alto Jacuí sub-basin due to the lack of extensive collecting effort in the region. The expansion of the sampled streams in Jacuí River basin may increase the records of species and information about endemic species.

ACKNOWLEDGEMENTS

We thank the colleagues of the Ichthyology Lab at the Universidade Federal do Rio Grande do Sul for field work; MSc. J. Ferrer, MSc. P.C. Silva for help in identifying the species studied and Dr. L.R. Malabarba to provide the species figures; the Orlandi and Bonato families for help and support in field work; and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Proc. 1104786, to KOB) for financial support. We also thank to Adam J. Taylor for reviewing the English of this manuscript.

LITERATURE CITED

- Agostinho, A.A., S.M. Thomaz and L.C. Gomes. 2005. Conservação da biodiversidade em águas continentais do Brasil. *Megadiversidade* 1(1): 70–78.
- Alves, T.P. and N.F. Fontoura. 2009. Statistical distribution models for migratory fish in Jacuí basin, South Brazil. *Neotropical Ichthyol-*

- ogy 7(4): 647–658. doi: [10.1590/S1679-62252009000400014](https://doi.org/10.1590/S1679-62252009000400014)
- Azpelicueta, M. de las M., J.G. Lundberg and M. Loureiro. 2008. *Pimelodus pintado* (Siluriformes: Pimelodidae), a new species of catfish from affluent rivers of Laguna Merín, Uruguay, South America. Proceedings of the Academy of Natural Sciences of Philadelphia 157: 149–162. doi: [10.1635/0097-3157\(2008\)157\[149:PPSPAN\]2.0.CO;2](https://doi.org/10.1635/0097-3157(2008)157[149:PPSPAN]2.0.CO;2)
- Barletta, M., A.J. Jaureguizar, C. Baiguns, N.F. Fontoura, A.A. Agostinho, V.M.F. Almeida-Val, A.L. Val, R.A. Torres, L.F. Jimenes-Segura, T. Giarrizzo, N.N. Fabré, V.S. Batista, C. Lasso, D.C. Taphorin, M.F. Costa, P.T. Chaves, J.P. Vieira and M.F.M. Corrêa. 2010. Fish and aquatic habitat conservation in South America: a continental overview with emphasis on neotropical systems. Journal of Fish Biology 76: 2118–2176. doi: [10.1007/s10641-014-0226-y](https://doi.org/10.1007/s10641-014-0226-y)
- Bastos, J.R.H. 2002. Biologia alimentar da taxocenose de peixes do rio Silveira (cabeceira do rio Pelotas), São José dos Ausentes, Rio Grande do Sul, Brasil [M.Sc. dissertation]. Porto Alegre: Universidade Federal do Rio Grande do Sul. 67 pp.
- Becker, F.G. 2002. Distribuição e abundância de peixes de corredeiras e suas relações com características de habitat local, bacia de drenagem e posição espacial em riachos de mata Atlântica (bacia do rio Maquine, RS, Brasil) [Ph.D. thesis]. São Paulo: Universidade Federal de São Carlos. 190 pp.
- Becker, F.G., L.C.C. De Fries, J. Ferrer, V.A. Bertaco, K.D.G. Luz-Agostinho, J.F.P. Silva, A.R. Cardoso, Z.M.S. Lucena and C.A.S. Lucena. 2013. Fishes of the Taquari-Antas river basin (Patos Lagoon basin), Southern Brazil. Brazilian Journal of Biology 73(1): 79–90. doi: [10.1590/S1519-69842013000100010](https://doi.org/10.1590/S1519-69842013000100010)
- Bertaco, A. and C.A.S. Lucena. 2010. Redescription of *Astyanax obscurus* (Hensel, 1870) and *A. laticeps* (Cope, 1894) (Teleostei: Characidae): two valid freshwater species originally described from rivers of Southern Brazil. Neotropical Ichthyology 8(1): 7–20. doi: [10.1590/S1679-62252010000100002](https://doi.org/10.1590/S1679-62252010000100002)
- Betancur, R.R., R.E. Broughton, E.O. Wiley, K. Carpenter, J.A. López, C. Li, N.I. Holcroft, D. Arcila, M. Sanciangco, J.C. Cureton, F. Zhang, T. Buser, M.A. Campbell, J.A. Balesteros, A. Roa-Varon, S. Willis, W.C. Borden, T. Roelwy, P.C. Reneau, D.J. Hough, G. Lu, T. Grande, G. Arratia and G. Ortí. 2013. The tree of life and a new classification of bony fishes. PLoS Currents. doi: [10.1371/currents.tol.53ba26640df0ccaee75bb165c8c26288](https://doi.org/10.1371/currents.tol.53ba26640df0ccaee75bb165c8c26288)
- Bloch, M.E. 1794. Naturgeschichte der ausländischen Fische. Berlin: Hesse. 8: 1–174. <http://biodiversitylibrary.org/page/41367521>
- Bloch, M.E. 1795. Naturgeschichte der ausländischen Fische. Berlin: Hesse. 9: 1–192. <http://biodiversitylibrary.org/page/41367701>
- Bockmann, F.A. 1998. Análise filogenética da família Heptapteridae (Teleostei, Ostariophysi, Siluriformes) e redefinição de seus gêneros [Ph.D. thesis]. São Paulo: Universidade de São Paulo. 599 pp.
- Boulenger, G.A. 1887. Descriptions of new South-American characineid fishes. Annals and Magazine of Natural History (5th series) 19(111): 172–174. <http://biodiversitylibrary.org/page/16334767>
- Bonato, K.O. and J. Ferrer. 2013. New record and distribution extension of *Phalloceros spiloura* Lucinda, 2008 (Cyprinodontiformes: Poeciliidae). Check List 9(6): 1545–1547. doi: [10.15560/9.6.1545](https://doi.org/10.15560/9.6.1545)
- Bozzetti, M. and U.H. Schultz. 2004. An index of biotic integrity base on fish assemblages for subtropical streams in southern Brazil. Hydrobiologia 529: 133–144. doi: [10.1007/s10750-004-5738-6](https://doi.org/10.1007/s10750-004-5738-6)
- Buckup, P.A. and R.E. Reis. 1997. Characidiin genus *Characidium* (Teleostei, Characiformes) in southern Brazil, with description of three new species. Copeia 1997(3): 531–548. doi: [10.2307/1447557](https://doi.org/10.2307/1447557)
- Buckup, P.A., N.A. Menezes and M.S. Ghazzi. 2007. Catálogo das Espécies de Peixes de Água Doce do Brasil. Rio de Janeiro: Museu Nacional, Universidade Federal do Rio de Janeiro, Série Livros 23. 195 pp.
- Carvalho, T.P., P. Lehmann, A.E.H.L. Pereira and R.E. Reis. 2008. A new species of *Hisonotus* (Siluriformes: Loricariidae: Hypoptopomatinae) from the Laguna dos Patos basin, southern Brazil. Copeia 2008(3): 510–516. doi: [10.1643/CI-07-130](https://doi.org/10.1643/CI-07-130)
- Carvalho, T.P. and R.E. Reis. 2011. Taxonomic review of *Hisonotus* Eigenmann & Eigenmann (Siluriformes: Loricariidae: Hypoptopomatinae) from the laguna dos Patos system, southern Brazil. Neotropical Ichthyology 9(1): 1–48. doi: [10.1590/S1679-62252011000100001](https://doi.org/10.1590/S1679-62252011000100001)
- Castro, R.M.C. 1999. Evolução da ictiofauna de riachos sul-americanos: padrões gerais e possíveis processos causais; pp. 139–155, in E.P. Caramaschi, R. Mazzoni and R.P. Peres-Neto (eds.). Ecologia de Peixes de Riachos: Estado Atual e Perspectivas. Rio de Janeiro: Série Oecologia Brasiliensis.
- COAJU (Comitê de gerenciamento de Bacias Hidrográficas do Alto Jacuí). [2009]. Plano de Gerenciamento da Bacia hidrográfica do Alto Jacuí. Relatório-T2. Consolidação das informações existentes sobre os recursos hídricos da bacia. Accessed at http://www.upf.br/coaju/index.php?option=com_content&task=view&id=288&Itemid=55, 20 June 2012
- Cope, E.D. 1894. On the fishes obtained by the Naturalist Expedition in Rio Grande do Sul. Proceedings of the American Philosophical Society 33: 84–108. <http://biodiversitylibrary.org/page/29851042>
- Costa, P.F. and U.H. Schulz. 2010. The fish community as an indicator of biotic integrity of the streams in the Sinos River basin, Brazil. Brazilian Journal of Biology 70(4): 1195–1205. doi: [10.1590/S1519-69842010000600009](https://doi.org/10.1590/S1519-69842010000600009)
- Dufech, A.P.S. and C.B. Fialho. 2006. Biologia populacional de *Pachyurus bonariensis* Steindachner, 1879 (Perciformes, Sciaenidae), uma espécie alóctone no sistema hidrográfico da laguna dos Patos, Brasil. Biota Neotropica 7(1): 105–110. <http://www.biotaneotropica.org.br/v7n1/pt/abstract?article+bn03207012007%20ISSN%201676-0603>
- Dufech, A.P.S. and C.B. Fialho. 2009. Estudo comparado da taxocenose de peixes em dois ambientes aquáticos do Parque Estadual de Itapuã, sul do Brasil. Iheringia, Série Zoologia 99(2): 177–188. doi: [10.1590/S0073-47212009000200010](https://doi.org/10.1590/S0073-47212009000200010)
- Eigenmann, C.H. and R.S. Eigenmann. 1888. Preliminary notes on South American Nematognathi. Proceedings of the California Academy of Sciences 1(2): 119–172. <http://biodiversitylibrary.org/page/7133491>
- Ferreira, C.P. and L. Casatti. 2006. Influência da estrutura do habitat sobre a ictiofauna de um riacho em uma micro-bacia de pastagem, São Paulo, Brasil. Revista Brasileira de Zoologia 23(3): 642–651. doi: [10.1590/S0101-81752006000300006](https://doi.org/10.1590/S0101-81752006000300006)
- Ferrer, J. and L.R. Malabarba. 2013. Taxonomic review of genus *Trichomycterus* Valenciennes (Siluriformes: Trichomycteridae) from the laguna dos Patos system, Southern Brazil. Neotropical Ichthyology 11(2): 217–246. doi: [10.1590/S1679-62252013000200001](https://doi.org/10.1590/S1679-62252013000200001)
- Fialho, C.B., L.C. Schifino and J.R. Verani. 1998. Biologia reprodutiva de *Oligosarcus jenynsii* (Günther) (Characiformes, Characidae) da Lagoa das Custódias, Tramandaí, Rio Grande do Sul, Brasil. Revista Brasileira de Zoologia 15(3): 775–782.
- Flores-Lopes, F., M. Cetra and L.R. Malabarba. 2010. Utilização de índices ecológicos em assembléias de peixes como instrumento de avaliação da degradação ambiental em programas de monitoramento. Biota Neotropica 10(4): 183–193. <http://www.biotaneotropica.org.br/v10n4/pt/abstract?article+bn03710042010>
- Froese, R. and D. Pauly. [2014]. FishBase. Accessed at <http://www.fishbase.org>, 20 October 2014.
- FEPAM (Fundação Estadual de Proteção Ambiental Henrique Luiz Roessler – RS). [2012]. Base de Dados. Accessed at <http://www.fepam.rs.gov.br/qualidade/guaiba.asp>, 25 June 2012.
- Galves, W., O.A. Shibatta, O.A. and F.C. Jerep. 2009. Estudos sobre a diversidade de peixes da bacia do alto rio Paraná: uma revisão

- histórica. *Semina: Ciência Biológicas e da Saúde* 30(2): 141–154.
- Garcia, A.M. and J.P. Vieira, J.P. 2001. O aumento da diversidade de peixes no estuário da lagoa dos Patos durante o episódio *El Niño* 1997–1998. *Atlântica* 23: 85–96.
- Garcia, A.M., M.B. Raseira, J.P. Vieira, K.O. Winemiller and A.M. Grimm. 2003. Spatiotemporal variation in shallow-water freshwater fish distribution and abundance in a large subtropical coastal lagoon. *Environmental Biology of Fishes* 68: 215–228. doi: [10.1023/A:1027366101945](https://doi.org/10.1023/A:1027366101945)
- Garcia, A.M., A.M. Bemvenuti, J.P. Vieira, D.M.L.M. Marques, M.D.M. Burns, A. Moresco and M.V.L. Condini. 2006. Checklist comparasion and dominance patterns of the fish fauna at Taim wetland, South Brazil. *Neotropical Ichthyology* 4(2): 261–268. doi: <http://dx.doi.org/10.1590/S1679-62252006000200012>
- Gomes, A.L. 1947. A small collection of fishes from Rio Grande do Sul, Brazil. *Miscellaneous Publications, Museum of Zoology, University of Michigan* 67: 1–39. <http://hdl.handle.net/2027.42/56312>
- Günther, A. 1864. Catalogue of the fishes in the British Museum. Catalogue of the Physostomi, containing the families Siluridae, Characinidae, Haplochitonidae, Sternoptychidae, Scopelidae, Stomiatidae in the collection of the British Museum 5: 1–455. <http://biodiversitylibrary.org/page/8404768>
- Heckel, J. J. 1840. Johann Natterer's neue Flussfische Brasilien's nach den Beobachtungen und Mittheilungen des Entdeckers beschrieben (Erste Abtheilung, Die Labroiden). *Annalen des Wiener Museums der Naturgeschichte* 2: 325–471. <http://biodiversitylibrary.org/item/127839>
- Hensel, R. 1868. Beiträge zur Kenntniss der Wirbelthiere Südbrasilien's. *Archiv für Naturgeschichte* 34: 323–375.
- Hensel, R. 1870. Beiträge zur Kenntniss der Wirbelthiere Südbrasilien's. *Archiv für Naturgeschichte* 36(1): 50–91. <http://biodiversitylibrary.org/page/7082229>
- Hirschmann, A. 2009. Composição e estrutura da assembleia de peixes em três arroios da sub-bacia do rio Forqueta, Sul do Brasil [M.Sc. dissertation]. Porto Alegre: Universidade Federal do Rio Grande do Sul. 70 pp.
- Langeani, F., R.M.C. Castro, O.T. Oyakawa, O.A. Shibatta, C.S. Pavanelli and L. Casatti. 2007. Diversidade da ictiofauna do Alto Rio Paraná: composição atual e perspectivas futuras. *Biota Neotropica* 7(3): 181–197. <http://www.biotaneotropica.org.br/v7n3/pt/abstract?article+bn03407032007>
- Linnaeus, C. 1758. *Systema Naturae*, Ed. X. (*Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Tomus I. Editio decima, reformata). *Holmiae* 1: 1–824. <http://biodiversitylibrary.org/item/10277>
- Lêvêque, C., T. Oberdorff, D. Paugy, M. Stiassny and P.A. Tedesco. 2008. Global diversity of fish (Pisces) in freshwater. *Hydrobiologia* 595: 545–567. doi: [10.1007/s10750-007-9034-0](https://doi.org/10.1007/s10750-007-9034-0)
- Lowe-McConnel, R.H. 1999. *Estudos Ecológicos de Comunidade de Peixes Tropicais*. São Paulo: EDUSP. 536 pp.
- Lucena, C.A.S., B.B. Calegari, E.H.L. Pereira and E. Dallegrave. 2013a. O uso de óleo de cravo na eutanásia de peixes. *Boletim Sociedade Brasileira de Ictiologia* 105: 20–24.
- Lucena, C.A.S., J.B. Castro and V.A. Bertaco. 2013b. Three new species of *Astyanax* from drainages of southern Brazil (Characiformes: Characidae). *Neotropical Ichthyology* 11(3): 537–552. doi: [10.1590/S1679-62252013000300007](https://doi.org/10.1590/S1679-62252013000300007)
- Lucena, C.A.S. and H.G. Soares. 2016. Review of species of the *Astyanax bimaculatus* “caudal peduncle spot” subgroup *sensu* Garutti & Langeani (Characiformes, Characidae) from the rio La Plata and rio São Francisco drainages and coastal systems of southern Brazil and Uruguay. *Zootaxa* 4072(1): 101–125. doi: [10.11646/zootaxa.4072.1.5](https://doi.org/10.11646/zootaxa.4072.1.5)
- Lucinda, P.H.F. 2008. Systematics and biogeography of the genus *Phalloceros* Eigenmann, 1907 (Cyprinodontiformes: Poeciliidae: Poeciliinae), with the description of twenty-one new species. *Neotropical Ichthyology* 6(2): 113–158. doi: [10.1590/S1679-62252008000200001](https://doi.org/10.1590/S1679-62252008000200001)
- Lütken, C.F. 1875. *Characinae novae Brasiliae centralis a clarissimo J. Reinhardt in provincia Minas-Geraes circa oppidulum Lagoa Santa in lacu ejusdem nominis, flumine Rio das Velhas et rivulis affluentibus collectae, secundum characteres essentialis breviter descriptae*. *Oversigt over det Kongelige Danske Videnskabernes Selskabs Forhandling og dets Medlemmers Arbejder*, Kjøbenhavn 1874(3): 127–143.
- Majolo, M.A. 2005. Estudo da taxocenose de peixes na Ilha do Andrade, bacia hidrográfica do rio Taquari, Arroio do Meio, Rio Grande do Sul – Brasil [M.Sc. dissertation]. Porto Alegre: Universidade Federal do Rio Grande do Sul. 120 pp.
- Malabarba, L.R. 1989. Histórico sistemático e lista comentada das espécies de peixes de água doce do sistema da Laguna dos Patos, Rio Grande do Sul, Brasil. *Comunicações do Museu de Ciências e Tecnologia da PUCRS, Série Zoologia* 2: 107–179.
- Malabarba, L.R. and S.H. Weitzman. 2003. Description of a new genus with six new species from southern Brazil, Uruguay and Argentina, with a discussion of a putative characid clade (Teleostei: Characiformes: Characidae). *Comunicações do Museu de Ciências e Tecnologia da PUCRS, Série Zoologia* 16(1): 67–151.
- Malabarba, L.R., C.B. Fialho, J.A. Anza J.A., J.F. Santos and G. N. Menezes. 2009. Peixes; pp. 131–156, in: I. Boldrini (ed.). *Biodiversidade dos Campos do Planalto das Araucárias*. Brasília: MMA.
- Maltchik, L., D.A. Müller and C. Stenert. 2005. Fish assemblages dynamics in a shallow floodplain lake in the South of Brazil. *Acta Limnologica Brasiliensia* 17(2): 185–198.
- Menezes, N.A. and A.C. Ribeiro. 2010. *Oligossarcus jacuiensis* (Characiformes: Characidae), a new species from the Uruguay and Jacuí River basins, southern Brazil. *Neotropical Ichthyology* 8(3): 649–653. doi: <http://dx.doi.org/10.1590/S1679-62252010000300010>
- Otoni, F.P. and M.M. Cheffe. 2009. A new species of *Australoheros* from the upper rio das Antas, laguna dos Patos System, southern Brazil. *Spixiana* 32(1): 153–159.
- Pelayo-Villamil, P., C. Guisande, P.R. Vari, A. Manjarés-Hernández, E. García-Rosello, J. González-Dacosta, J. Heine, L.G. Vilas, B. Patti, E.M. Quinci, L.F. Jiménez, C. Grnanado-Lorencio, P.A. Tedesco and J.M. Lobo. 2014. Global diversity patterns of freshwater fishes — potential victims of their own success. *Diversity and Distribution*. doi: [10.1111/ddi.12271](https://doi.org/10.1111/ddi.12271)
- Petry, A.C. and U.H. Schulz. 2006. Longitudinal changes and indicator species of the fish fauna in the subtropical Sinos River, Brazil. *Journal of Fish Biology* 69: 272–290. doi: [10.1111/j.1095-8649.2006.01110.x](https://doi.org/10.1111/j.1095-8649.2006.01110.x)
- Quoy, J.R.C. and J.P. Gaimard. 1824–1825. Description des Poissons; Chapitre IX, [1824]: 1–328, [1825]: 329–616, in: L. de Freycinet (ed.). *Voyage autour du monde: entrepris par ordre du roi ... exécuté sur les corvettes de S. M. l'Uranie et la Physicienne, pendant les années 1817, 1818, 1819 et 1820* Paris: Pillet. <http://biodiversitylibrary.org/page/40871183>
- Reis, R.E. and S.A. Schaefer. 1998. New cascudinhos from southern Brazil: systematics, endemism, and relationships (Siluriformes, Loricariidae, Hypoptopomatinae). *American Museum Novitates* 3254: 1–25. <http://hdl.handle.net/2246/3154>
- Reis, R.E., S.O. Kullander and C.J. Ferraris Jr. 2003. Check list of the freshwater fishes of South and Central America. Porto Alegre: EDIPUCRS. 742 pp.
- Regan, C.T. 1904. A monograph of the fishes of the family Loricariidae. *Transactions of the Zoological Society of London* 17(1): 191–350. <http://biodiversitylibrary.org/item/48598>
- Ribeiro, M.F. and A. Köhler. 2007. Ictiofauna do rio Pardinhas, RS, Brasil. *Cadernos de Pesquisa, Série Biologia* 19(2): 37–45.
- Rincán, O. and Kullander, S.O. 2008. The *Australoheros* (Teleostei:

- Cichlidae) species of the Uruguay and Paraná River drainages. *Zootaxa* 1724: 1–51.
- Rodriguez, M.S. and R.E. Reis, R.E. 2008. Taxonomic review of *Rineloricaria* (Loricariidae: Loricariinae) from the Laguna dos Patos drainage, southern Brazil, with the descriptions of two new species and the recognition of two species groups. *Copeia* 2008(2): 333–349. doi: [10.1643/CI-06-218](https://doi.org/10.1643/CI-06-218)
- Saccol-Pereira, A. and C.B. Fialho. 2010. Seasonal and diel variation in the fish assemblage of a Neotropical delta in southern Brazil. *Iheringia, Série Zoologia* 100(2): 169–178. doi: [10.1590/S0073-47212010000200013](https://doi.org/10.1590/S0073-47212010000200013)
- SEMA (Secretaria do Meio Ambiente). Base de dados. Accessed at http://www.sema.rs.gov.br/conteudo.asp?cod_menu=56&cod_conteudo=6250, 20 June 2012.
- Silva, J.F.P. 1998. Revisão taxonômica das espécies de *Bryconamericus* (Eigenmann, 1907) do sul e sudeste do Brasil (Ostariophysi, Characidae) [M.Sc. dissertation]. Porto Alegre: Pontifícia Universidade Católica do Rio Grande do Sul. 107 pp.
- Silva, G.N.M. 2009. Convergência ecomorfológica entre comunidades de peixes com histórias biogeográficas distintas [M.Sc. dissertation]. Porto Alegre: Universidade Federal do Rio Grande do Sul. 47 pp.
- Súarez, Y.R. and M. Petrere-Júnior. 2005. Organização das assembleias de peixes em riachos da bacia do rio Iguatemi, Estado do Mato Grosso do Sul. *Acta Scientiarum* 27(2): 161–167. doi: [10.4025/actasciobiolsci.v27i2.1350](https://doi.org/10.4025/actasciobiolsci.v27i2.1350)
- Tagliani, P.R.A. 1994. Ecologia da assembleia de peixes de três riachos da planície costeira do Rio Grande do Sul. *Atlântica* 16: 55–68.
- Thomaz, A.T., D. Arcila, D., G. Ortí and L.R. Malabarba. 2015. Molecular phylogeny of the subfamily Stevardiinae Gill, 1858 (Characiformes: Characidae) classification and the evolution of reproductive traits. *BMC Evolutionary Biology* 15: 146. doi: [10.1186/s12862-015-0403-4](https://doi.org/10.1186/s12862-015-0403-4)
- Valenciennes, A. 1834–1842. Poissons; pp. 1–16, in: A. d'Orbigny (ed.). *Voyage dans l'Amérique méridionale: (le Brésil, la république orientale de l'Uruguay, la République argentine, la Patagonie, la république du Chili, la république de Bolivie, la république du Pérou)*, exécuté pendant les années 1826, 1827, 1828, 1829, 1830, 1831, 1832, et 1833.
- Vannote, R.L., G.W. Mishall, K.W. Cummins, J.R. Sedell and C.E. Cushing. 1980. The River Continuum Concept. *Canadian Journal of Fisheries and Aquatic Sciences* 37(1): 130–137. doi: [10.1139/f80-017](https://doi.org/10.1139/f80-017)
- Vidal, L.V.O., R.C.B. Albinati, A.C.L. Albinati, A.D. Lira, T.R. Almeida and G.B. Santos. 2008. Eugenol como anestésico para a tilápia do Nilo. *Pesquisa Agropecuária Brasileira* 43(8): 1069–1074.
- Vilella, F.S., F.G. Becker, S.M. Hartz and G. Barbier. 2004. Relation between environmental variables and aquatic megafauna in a first order stream of the Atlantic Forest, southern Brazil. *Hydrobiologia* 528: 17–30. doi: [10.1007/s10750-004-0688-6](https://doi.org/10.1007/s10750-004-0688-6)
- Volcan, M.V., A.C. Gonçalves and L.E.K. Lanés. 2011. Distribution, habitat and conservation status of two threatened annual fishes (Rivulidae) from southern Brazil. *Endangered Species Research* 13: 79–85. doi: [10.3354/esr00316](https://doi.org/10.3354/esr00316)
- Wiley, E.O. and G.D. Johnson. 2010. A teleost classification based on monophyletic groups; pp. 123–182, in: J.S. Nelson, H.P. Schultze and M.V.H. Wilson (eds.). *Origin and Phylogenetic Interrelationships of Teleosts*. München: Verlag Dr. Friedrich Pfeil.
- Winckler-Sosinski, L.T., A. Schwarzbald and U.H. Schulz. 2009. Fish assemblage structure in altitude rivers under the effect of exotic species introduction, northeast of Rio Grande do Sul, Brazil. *Acta Limnologica Brasiliensia* 21(4): 473–482.
- Winemiller, K.O., A.A. Agostinho and P.E. Caramaschi. 2008. Fish ecology in tropical streams; pp. 336–346, in: D. Dudgeon (ed.). *Tropical Stream Ecology*. San Diego, California: Academic Press.

Author contributions: KOB collected data, verified the specimens and wrote the text. CBF wrote the text.

Received: 17 November 2015

Accepted: 17 March 2016

Academic editor: Tiago P. Carvalho